

Minimal measurements of the gate fidelity of a qudit map

E. Bagan, M. Baig and R. Muñoz-Tapia

*Grup de Física Teòrica & IFAE, Facultat de Ciències, Edifici Cn,
Universitat Autònoma de Barcelona, 08193 Bellaterra (Barcelona) Spain*

(Dated: July 26, 2002)

We obtain a simple formula for the average gate fidelity of a linear map acting on qudits. It is given in terms of minimal sets of pure state preparations alone, which may be interesting from the experimental point of view. These preparations can be seen as the outcomes of certain minimal positive operator valued measures. The connection of our results with these generalized measurements is briefly discussed.

PACS numbers: 03.65.Bz, 03.67.-a

The interaction with the environment has a significant negative impact on any physical implementation of a unitary gate or a quantum channel. Decoherence, turning pure states into mixed states, creeps up and unitarity is lost in the Hilbert subspace of the signal states. The characterization of the quality of such implementations is, hence, of utmost importance for quantum computation from both experimental and theoretical point of view [1].

A physical quantum gate or channel is best described by a trace-preserving linear map or superoperator $\mathcal{E}$, which is assumed to be an approximation of a unitary operator U characterizing the quantum gate ($U = \mathbb{1}$ for a quantum channel). The average gate fidelity

$$\overline{F}(\mathcal{E}, U) = \int d\psi \operatorname{tr} [U \rho_\psi U^\dagger \mathcal{E}(\rho_\psi)], \quad (1)$$

where $d\psi$ is the invariant Haar measure on the space of pure states $|\psi\rangle$ (i.e., $d\psi = d\psi'$ if $|\psi'\rangle = U|\psi\rangle$) and $\rho_\psi = |\psi\rangle\langle\psi|$, is recognized as a convenient figure of merit and is widely used to assess the quality of a quantum gate or channel.

For gates acting on qubits, Bowdrey *et al.* [2] have recently derived a simple formula for $\overline{F}(\mathcal{E}, U)$, from which they obtain a convenient way of actually measuring the gate fidelity in a laboratory. It amounts to replacing the integral in (1) by a sum over a finite number of pure states whose Bloch vectors point at the vertices of an octahedron or a tetrahedron inscribed in the Bloch sphere. Nielsen [3] has obtained a similar formula for qudits (i. e., quantum states belonging to a d -dimensional Hilbert space) in terms of a finite set of unitary operations U_j orthogonal with respect to the Hilbert-Schmidt inner product, namely, such that $\operatorname{tr} (U_i^\dagger U_j) = d \delta_{ij}$. His derivation is based on Horodeckis' formula connecting $\overline{F}(\mathcal{E}, \mathbb{1}) \equiv \overline{F}(\mathcal{E})$ with the entanglement fidelity [4], for which he also provides a simple proof. Our aim is to give a more natural generalization of Bowdrey's formula so that only measurements over a (minimum) number of pure states need to be performed to verify $\overline{F}$ experimentally. Interestingly enough, the solution of this problem is not far removed from that of obtaining minimal positive operator valued measurements (POVMs) in the context

of optimal communication of directions through a quantum (spin) channel, which has received much attention over the last few years [5, 6, 7].

In the first part of this letter we derive a general expression of $\overline{F}$ in terms of the $SU(d)$ group generators. In the second part we write $\overline{F}$ as an average of measurements over a finite and minimal number of pure state preparations, which may be experimentally relevant.

We first notice that due to the invariance of $d\psi$ one has $\overline{F}(\mathcal{E}, U) = \overline{F}(\mathcal{E}', \mathbb{1}) = \overline{F}(\mathcal{E}')$, where $\mathcal{E}'(\rho) \equiv \mathcal{E}(U^\dagger \rho U)$. Hence we only need to consider the simpler form $\overline{F}(\mathcal{E})$ without any loss of generality. Furthermore, the uniform measure $d\psi$ can be effectively realized in the following way: $\int d\psi \mathcal{F}(\psi) \equiv \int dU \mathcal{F}(U\psi_0)$, where $\mathcal{F}$ is any function of $|\psi\rangle$, $|\psi_0\rangle$ is a fixed reference state, and dU is the Haar measure of $SU(d)$ normalized such that $\int dU = 1$. This ensures that $\int d\psi \mathcal{F}(\psi) = \int d\psi \mathcal{F}(U'\psi)$ for any $U' \in SU(d)$. Obviously, not all the $d^2 - 1$ parameters involved in dU are physically significant. E.g., for qubits ($d = 2$), $d\psi = dn$ is the uniform measure on the 2-sphere S^2 , which can be parametrized by the Euler angles α and β . If $|\psi_0\rangle$ is an eigenstate of the Pauli matrix σ_z , any function of $|\psi\rangle$ is independent of the third Euler angle γ . Hence, $\int d\psi \mathcal{F} = [\int d\gamma / (2\pi) = 1] \times \int dn \mathcal{F} = \int dU \mathcal{F}$. For qudits, dn must be replaced by the invariant measure of $SU(d)/[SU(d-1) \times U(1)]$, since any reference state $|\psi_0\rangle$ is now invariant under $SU(d-1) \times U(1)$. Note that the number of parameters match, as qudits depend on $2(d-1)$ real variables. With all this we finally have

$$\overline{F}(\mathcal{E}) = \int dU \operatorname{tr} [U \rho_0 U^\dagger \mathcal{E}(U \rho_0 U^\dagger)], \quad (2)$$

where we have defined $\rho_0 \equiv |\psi_0\rangle\langle\psi_0|$.

There exists a $(d^2 - 1)$ -dimensional unitary vector with components n_a^0 , $a = 1, 2, \dots, d^2 - 1$, such that the density matrix ρ_0 can be written as

$$\rho_0 = \frac{\mathbb{1}}{d} + k_d n_0^a T_a \equiv \frac{\mathbb{1}}{d} + k_d \vec{n}_0 \cdot \vec{T}, \quad (3)$$

where $k_d = \sqrt{2(d-1)/d}$ and $\{T_a\}$ are the (hermitian and traceless) generators of $SU(d)$, normalized so that $\operatorname{tr} (T_a T_b) = \delta_{ab}/2$. They can be chosen as $T_a = \lambda_a/2$,

where λ_a are a generalization of the Gell-Mann matrices of $SU(3)$. Throughout this letter a sum over repeated latin indexes is understood. It is straightforward to obtain

$$\vec{n}_0 = \frac{2}{k_d} \text{tr}(\rho_0 \vec{T}). \quad (4)$$

We now recall the well known relation

$$UT_a U^\dagger = (\text{Ad}U)_a^b T_b, \quad (5)$$

where Ad stands for the adjoint representation of $SU(d)$. We also recall the orthogonality of the irreducible representations of compact groups, which in the present case implies

$$\int dU (\text{Ad}U)_a^b = 0, \quad (6)$$

$$\int dU (\text{Ad}U)_a^b (\text{Ad}U)_c^d = \frac{\delta_{ac} \delta^{bd}}{d^2 - 1}. \quad (7)$$

Using that $\mathcal{E}$ is linear and trace-preserving one gets

$$\overline{F}(\mathcal{E}) = \frac{1}{d} + \frac{2}{d(d+1)} \sum_{a=1}^{d^2-1} \text{tr} [T_a \mathcal{E}(T_a)]. \quad (8)$$

This is the generalization of Bowdrey's formula

$$\overline{F}(\mathcal{E}) = \frac{1}{2} + \frac{1}{3} \sum_{i=x,y,z} \text{tr} \left[\frac{\sigma_i}{2} \mathcal{E} \left(\frac{\sigma_i}{2} \right) \right] \quad (9)$$

for qubits. That concludes the first part of this letter.

The key ingredient of the derivation above is the orthogonality relation of the group representations, exemplified by Eqs. 6 and 7. We will show that it is possible to find a discrete version of these equations. Namely, one can find a *finite* set of $SU(d)$ elements $\{U_r\}$ and positive constants $\{c_r\}$ such that

$$\sum_r c_r (\text{Ad}U_r)_a^b = 0 \quad (10)$$

$$\sum_r c_r (\text{Ad}U_r)_a^b (\text{Ad}U_r)_c^d = \frac{\delta_{ac} \delta^{bd}}{d^2 - 1}. \quad (11)$$

With this we can reverse the steps going from (2) to (8) using the relations (10) and (11) instead of their continuous version (6) and (7), and obtain

$$\overline{F}(\mathcal{E}) = \sum_r c_r \text{tr} [\rho_r \mathcal{E}(\rho_r)]. \quad (12)$$

This equation has a very convenient form which allows setting up experimental tests to determine the fidelity of a gate or channel, as will be discussed below.

Let us briefly discuss the solutions to Eqs. 10 and 11. The idea is to generalize the concept of a finite set of *isotropically distributed* unit vectors introduced in [6] and

adapt it to the problem at hand. A sufficient condition for (12) to hold can be obtained by contracting (10) and (11) with n_0^a . If we define

$$n_0^a (\text{Ad}U_r)_a^b \equiv n_r^b, \quad (13)$$

conditions (10) and (11) lead to

$$\sum_r c_r n_r^b = 0; \quad \sum_r c_r n_r^b n_r^d = \frac{\delta^{bd}}{d^2 - 1}. \quad (14)$$

In this sense, we may qualify the set $\{n_r\}$ as isotropically distributed (as far as the adjoint representation $SU(d)$ is concerned). We have traded the problem of finding $\{U_r\}$ for that of finding $\{n_r\}$. Note, however that the set of matrices $\{\text{Ad}U\}$ is a proper subgroup of $SO(d^2 - 1)$. Hence, not any vector on the sphere $\mathbb{S}^{d^2-2}$ is admissible. We will come back to this issue below. The relations (10), (11) and (14) also appear in the rather different context: the construction of finite positive operator valued measurements which are optimal for communicating a direction [6, 7, 8]. Leaving aside an overall trivial normalization (from Eq. 14 it follows that $\sum_r c_r = 1$ whereas in [8] $\sum_r c_r = d$), the results in those papers can be readily used here. In particular, it is proved in [8] that solutions of (14) exist and the minimal one is given by a set of d^2 vectors pointing at the vertices of a regular hypertetrahedron or, more properly, $(d^2 - 1)$ -simplex, inscribed on $\mathbb{S}^{d^2-2}$. This hypertetrahedron is defined by the condition

$$\vec{n}_r \cdot \vec{n}_s = -\frac{1}{d^2 - 1}, \quad r \neq s, \quad (15)$$

and the exact overall orientation has to be chosen so that all vectors $\vec{n}_r$ are of the form (13). For this hypertetrahedron all the coefficients c_r are equal: in our notation $c_r = 1/d^2$, $r = 1, 2, \dots, d^2$. A explicit form of $\vec{n}_r$ for $SU(3)$ can be found in [8], where also the general case is briefly discussed. The solution is more conveniently expressed in terms of states $|\psi_r\rangle$ such that

$$|\psi_r\rangle\langle\psi_r| \equiv \rho_r = U_r \rho_0 U_r^\dagger = \frac{\mathbb{1}}{d} + k_d \vec{n}_r \cdot \vec{T}. \quad (16)$$

Then (15) translates into

$$|\langle\psi_r|\psi_s\rangle|^2 = \frac{1}{d+1}, \quad r \neq s. \quad (17)$$

Since this equation is a condition on states, Eq 13 is automatically satisfied by the corresponding Bloch vectors.

For $SU(3)$ a solution of (17) has the simple form

$$|\psi_r\rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ e^{2(r-1)\pi i/3} \\ 0 \end{pmatrix}, \quad r = 1, 2, 3. \quad (18)$$

The remaining six states are obtained by applying cyclic permutations to the components of $|\psi_{1-3}\rangle$ (a different

choice of states is given in [8]). From an experimental point of view, the states (18) have a very appealing form; each of them involves a linear combination of *only two* states of the computational basis. If the qutrit is implemented by say three atomic levels, only two levels need to be manipulated to prepare each one of the ρ_r .

In the $SU(2)$ case, a solution in terms of $\vec{n}_r$ is more simple and transparent, mainly because $SU(2)$ is isomorphic to $SO(3)$, which makes (13) trivially satisfied by any vector of S^2 . A compact solution is given by

$$\vec{n}_1 = \frac{1}{\sqrt{3}}(1, 1, 1), \quad \vec{n}_2 = \frac{1}{\sqrt{3}}(-1, -1, 1), \quad (19)$$

and $\vec{n}_3, \vec{n}_4$ are again obtained by applying cyclic permutations to the components of $\vec{n}_2$.

For the hypertetrahedra discussed above, Eq. 12 can be cast as

$$\overline{F}(\mathcal{E}) = \frac{1}{d^2} \sum_{r=1}^{d^2} \text{tr} [\rho_r \mathcal{E}(\rho_r)], \quad (20)$$

which is the $SU(d)$ generalization of Bowdrey's $SU(2)$ formula. This equation is our main result and provides a remarkably simple procedure for measuring $\overline{F}(\mathcal{E})$. One just has to average the fidelities for d^2 isotropically distributed pure states. Notice that all state preparations ρ_r have the same weight in the average, thus reducing systematic errors. In some sense, Eq. 20 can be regarded as the average survival rate of the states $\{\rho_r\}$ in a quantum channel characterized by the linear map $\mathcal{E}$. There is an alternative way of writing (20) which may provide further insight. It is straightforward to verify that the projectors $O_r \equiv \rho_r/d$ are the complete set of positive operators (i.e., $\sum O_r = \mathbb{1}$) of a minimal POVM. Thus, the states ρ_r are just the preparations produced by the device characterized by $\{O_r\}$. We can write (20) as

$$\overline{F}(\mathcal{E}) = \frac{1}{d} \sum_{r=1}^{d^2} \text{tr} [O_r \mathcal{E}(\rho_r)]. \quad (21)$$

We readily see that the same device $\{O_r\}$ could be used for the preparations of the pure states $\{\rho_r\}$ as well as for the measurements over $\{\mathcal{E}(\rho_r)\}$.

Although we have presented the results for the minimal sets of ρ_r , for practical reasons, one may wish to use a larger number of states. This possibility is easily implemented in this framework, as conditions (13) and (14) are not specific of minimal sets but entirely general. E.g., for qubits one can find a set of six states whose Bloch vectors point at the vertices of a regular octahedron [2, 5, 7]). In this case there is a simple setting for preparing the states, just three Stern-Gerlach's oriented along the three orthogonal directions.

We are grateful to A. Acín and A. Bramon for helpful conversations. Financial support from contracts CI-CYT AEN99-0766 and CIRIT 2000SGR-00063 is acknowledged.

-
- [1] J.F. Poyatos, J.I. Cirac and P. Zoller, Phys. Rev. Lett. **78**, 390 (1995).
 - [2] M.D. Bowdrey *et al.*, Phys. Lett. **A294** 258 (2002).
 - [3] M.A. Nielsen, arXiv:quant-ph/0205035.
 - [4] M. Horodecki, P. Horodecki and R. Horodecki, Phys. Rev. **A60** 1888 (1999).
 - [5] S. Massar and S. Popescu, Phys. Rev. Lett. **74**, 1259 (1995); N. Gisin and S. Popescu, Phys. Rev. Lett. **83**, 432 (1999); E. Bagan *et al.*, Phys. Rev. Lett. **85**, 5230 (2000); A. Peres and P. F. Scudo, Phys. Rev. Lett. **86**, 4160 (2001).
 - [6] E. Bagan, M. Baig and R. Muñoz-Tapia, Phys. Rev. **A64** 022305 (2001).
 - [7] J.I. Latorre, P. Pascual and R. Tarrach Phys. Rev. Lett. **81** 1351 (1998).
 - [8] A. Acín, J.I. Latorre and P. Pascual, Phys. Rev. **A61** 022113 (2000).